

Cardboard floor: About the barriers for social progression and their impact on the representativeness of epidemiological studies.

Jorge Arias de la Torre, MSc PhD^{1,2}, Jose M. Valderas, MD PhD³, Fernando G. Benavides, MD PhD^{2,4}, Jordi Alonso MD PhD^{2,4,5}.

¹ Institute of Psychiatry, Psychology and Neurosciences (IoPPN), King's College London, London, UK.

² CIBER Epidemiology and Public Health (CIBERESP), Madrid, Spain.

³ Health Services and Policy Research Group, University of Exeter Medical School, Exeter, UK.

⁴ Dept. of Experimental and Health Sciences, Pompeu Fabra University (UPF), Barcelona, Spain.

⁵ Health Services Research Group, Hospital del Mar Medical Research Institute (IMIM), Barcelona, Spain.

Word count: 846

Corresponding author:

Jorge Arias de la Torre

Department of Psychological Medicine, Division of Academic Psychiatry

Institute of Psychiatry, Psychology and Neuroscience (IoPPN)

King's College London, De Crespigny Park, London SE5 8AF

+44 754309727 | Jorge.arias_de_la_torre@kcl.ac.uk

23 **Abstract (111 words)**

24 The most disadvantaged extreme of the social continuum is usually underrepresented in
25 epidemiological studies. We discuss the consequences of excluding this segment of the
26 population and suggest different approaches for addressing this issue. In particular, we
27 describe/analyse a barrier that tends to perpetuates people in the most disadvantaged
28 extreme of the social continuum, hereinafter referred to as the “cardboard floor”. Besides,
29 we propose different approaches to accessing to the least favoured, segment in order to
30 study the cardboard floor. The adoption of these strategies could help to visualize this
31 barrier, allowing to better monitoring social mobility and their expected health
32 improvements, as well as increasing the representativity of population health studies.

33

A continuum of socioeconomic status ranging from the least to the most privileged persons is evidenced in population studies, with profound implications for health and care [1]. Individuals in the most disadvantaged social group suffer from extreme poverty and face several specific challenges to their health and health care [2]. They frequently cannot meet their most basic needs (including their physiological needs, most acutely exemplified by homelessness) and are at a higher risk of health problems and accelerated aging due to unhealthy habits (e.g. unhealthy diet and drug consumption), harmful environmental and biological factors, and social isolation [1–4]. As a result, the most socially disadvantaged persons have higher rates of premature mortality, especially caused by suicide and violence, and higher prevalence of all types of diseases, particularly infectious diseases and mental disorders [2,5]. Besides, care for chronic conditions is compromised for this population group, which relies to a substantial degree on emergency care, particularly in health systems that do not guarantee universal health coverage [5].

Even considering the relative size of the most deprived extreme of the social continuum (e.g. about 0.5% of the U.K. adult population in 2018 was considered homeless) [6], the scale of unmet health and health care needs would imply that improving their social mobility might have a significant impact on the overall health status of the population. However, several barriers significantly hinder this upward mobility. If a glass ceiling is used as a metaphor for the barrier to higher achievement, success, or recognition for individuals of certain groups within different careers or industries (e.g. women becoming CEOs), an even more appropriate one in this case would be a *cardboard floor*, making reference to the surface that is a daily experience for many extremely deprived people [7]. Studying the impact of this barrier on health, could help to understand it better, hopefully favouring social mobility. Conducting such studies, however, is not exempt from

difficulties, one which being particularly relevant: the lack of access to data from persons in the least favoured extreme of the social continuum.

The most disadvantaged group is very unlikely to be included in research and, as a result, is usually inadequately represented in health studies. This recruitment bias has important implications [3,4,8]. It limits the representativeness and external validity of surveys and population health studies and, furthermore, results in underestimation of the health risks, morbidity and mortality across the entire population. Importantly, it also hides the true scope of the specific issues affecting this group from researchers, policy makers and the public.

Different approaches focusing on improved sampling strategies to guarantee the representation of this group in population studies could be used. [3,8,9]. Proposed complementary strategies include: assigning greater sampling weights to individuals in this group, targeted over-recruitment, and/or intensifying fieldwork in marginal areas or suburbs through involvement of social organisations at local level. Nevertheless, these methods require some a priori knowledge of the number of people in this situation when defining the reference population for a specific study.

The use of data from administrative data and Electronic Health Records (EHR), such as the Medicaid claims data in the U.S. and the Clinical Practice Research Datalink (CPRD) in the U.K. [9,10], could also be a suitable way to access to the most socially disadvantaged persons. Relevant health and healthcare information for this population are often registered within these data sources. Some limitation of this data needs to be acknowledged in relation to their completeness, and ability to capture circumstances of maximum vulnerability and the inclusion of information on key mediating mechanisms relevant to determine biological, behavioural, and psychosocial pathways. However, such data also have strengths: they are in many cases mandatory, population-wide and usually

contain relevant information on different health outcomes, such as mortality or hospital admissions. Besides, most such data are potentially linkable to other relevant datasets for the study of this population (e.g. social care or demographic records) bringing together their strengths and, in some cases, allowing to overcome the abovementioned limitations [9,11]. Hence, the use of linked data from EHR could be a suitable way to capture relevant aspects of the most socially vulnerable individuals and, furthermore, might represent an adequate approach to obtain valid and reliable estimations of the health status in this part of the population. In addition linked EHR data, would allow estimating the relative numbers of the most disadvantaged group, providing relevant additional information on morbidity and outcomes and facilitating the implementation of improved sampling strategies [9].

The access to the most disadvantaged extreme segment of the social continuum remains a challenge for population health studies. Using a combination of approaches based on the use of HER linked data and strengthening the sampling strategy for the specific studies, might be a synergistic way to improve the validity of population health estimations. The adoption of these strategies could help to visualize the barriers for social mobility and the access to the most disadvantaged social groups. This will help to better understand the phenomena that perpetuate the cardboard floor and to shape care systems that truly “do not leave one behind” [12].

Contributorship Statement

All authors (JA-T, JMV, FGV and JA) were involved in all phases of the development of this manuscript, from the initial idea to the review and acceptance of the final draft for submission.

Competing interests

None declared.

Acknowledgements

We acknowledge Medical Research Council for its funding.

Funding

The present work is partially funded by the Medical Research Council (MR/S028188/1).

This article represents independent research part funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

References

- 1 World Health Organization. Social Determinants of Health: Key Concepts. *Who* 2014.
- 2 Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet* 2014;**384**:1529–40. doi:10.1016/S0140-6736(14)61132-6
- 3 Burnam MA, Koegel P. Methodology for Obtaining a Representative Sample of Homeless Persons. *Eval Rev* 1988;**12**:117–52. doi:10.1177/0193841X8801200202
- 4 Pizzi C, De Stavola B, Merletti F, *et al.* Sample selection and validity of exposure-disease association estimates in cohort studies. *J Epidemiol Community Heal* 2011;**65**:407–11. doi:10.1136/jech.2009.107185

- 5 Koh HK, O'Connell JJ. Improving health care for homeless people. JAMA - J.
Am. Med. Assoc. 2016. doi:10.1001/jama.2016.18760
- 6 Ministry of Housing Communities & Local Government. Live tables on
homelessness. Homelessness Stat. 2018.
- 7 Baldacci E, de Mello L, Inchauste G. Financial crises, poverty, and income
distribution. In: *Macroeconomic Policies and Poverty Reduction*. 2005.
doi:10.4324/9780203005804
- 8 Toro PA, Wolfe SM, Bellavia CW, *et al*. Obtaining representative samples of
homeless persons: A two-city study. *J Community Psychol* 1999;**27**:157–77.
doi:10.1002/(SICI)1520-6629(199903)27:2<157::AID-JCOP4>3.0.CO;2-2
- 9 Vickery KD, Shippee ND, Bodurtha P, *et al*. Identifying Homeless Medicaid
Enrollees Using Enrollment Addresses. *Health Serv Res* 2018;**53**:1992–2004.
doi:10.1111/1475-6773.12738
- 10 Herrett E, Gallagher AM, Bhaskaran K, *et al*. Data Resource Profile: Clinical
Practice Research Datalink (CPRD). *Int J Epidemiol* Published Online First:
2015. doi:10.1093/ije/dyv098
- 11 Padmanabhan S, Carty L, Cameron E, *et al*. Approach to record linkage of
primary care data from Clinical Practice Research Datalink to other health-related
patient data: overview and implications. *Eur J Epidemiol* Published Online First:
2019. doi:10.1007/s10654-018-0442-4
- 12 Griggs D, Stafford-Smith M, Gaffney O, *et al*. Policy: Sustainable development
goals for people and planet. *Nature*. 2013. doi:10.1038/495305a